

WORTHINGTON Stephen (1991) Karst
Hydrogeology of the Canadian Rocky
Mountains. dans Mémoire de Doctor of
Philosophy (Geography) (soutenu en mai 1991) ;
School of Graduate Studies - McMaster
University (Ontario, Canada) [Extrait : Contents
p.vii/xi - Figures p.xiii/xx - Tables p.xxi/xxiii]

Contents

List of figures

xiii

List of tables

xxi

Symbols used

xxv

Glossary

xxvii

Chapter 1 Introduction

1.1	Research objectives	1
1.2	Choice of study area	5
1.3	Organisation of the thesis	9
1.4	The problem of sampling bias	10

Chapter 2 The study area: Crowsnest Pass

2.1	Geology	13
2.1.1	Stratigraphy and mineralogy	13
2.1.2	Structure	18
2.2	Topography and drainage	20
2.2.1	Surface flow and inputs to the karst aquifers	20
2.2.2	Springs	22
2.3	Modern hydrogeology	27
2.3.1	Hydraulic conductivity of the karst aquifers	27
2.3.2	Groundwater flow vectors	29
2.4	Paleohydrogeology	30

Chapter 3 Hydrological and meteorological measurements

3.1	Introduction	31
3.2	Discharge measurement	32
3.3	Fluorometry	41
3.3.1	Aquifer hydraulics	41
3.3.2	Catchment delineation	52
3.4	Hydrochemistry	53
3.4.1	Specific conductivity and titrations	53
	Sampling and analysis	53
	The determination of aggressiveness	57
3.4.2	Neutron activation analysis	64
3.4.3	Stable isotope analysis	65
3.5	Meteorology	68
3.6	Characterisation of the springs at Crowsnest Pass	69

Chapter 4 The influence of boundary conditions on spring discharge and chemistry

4.1	Introduction	73
4.2	Underflow, full flow and overflow regimes	73
4.3	Springs of the Dinaric karst, Yugoslavia	81
4.4	Valles - San Luis Potosí region, Mexico	84
4.5	The Mendip Hills, England	88
4.6	Karst springs of Nittany Valley, Pennsylvania, USA	89
4.7	Karst springs of Ariège, France	94
4.8	Conclusion	98

Chapter 5 The hydraulics of karstic groundwater flow

5.1	Introduction	99
5.2	Diffuse flow and conduit flow	100
5.3	The Darcy-Weisbach equation	103
5.3.1	Velocity	103
5.3.2	Discharge	114
5.3.3	Area	115
5.3.4	Hydraulic gradient	117
5.3.5	Friction factor	124
5.4	Conclusion	126

Chapter 6 Flow nets in karst aquifers: morphometry and initial conditions

6.1	Introduction	127
6.2	Morphometric analysis of flow paths	130
6.2.1	Introduction	130
6.2.2	Conduit density	131
6.2.3	Conduit porosity	134
6.2.4	Segment length	135
6.2.5	Sinuosity	135
6.2.6	Flow belt width	140
6.2.7	Depth of loop crests and bases	143
6.2.8	Conclusion	149
6.3	Geological controls	149
6.3.1	Fracture occurrence	149
6.3.2	Primary tube development on multiple bedding planes	151
6.3.3	To what extent are bedding planes favoured?	155
6.4	Hydraulic controls on conduit development	158
6.4.1	Flow through fractures	158
6.4.2	Aquifer inputs	159
6.4.3	Initial hydraulic gradients	161
6.5	Topological controls	164
6.6	Kinetic controls on solution	166
6.7	Models of conduit development	169

Chapter 7 Flow nets in karst aquifers: the locus of flow during one generation

7.1	Introduction	173
7.2	Flow depth as a function of catchment length, dip and strike	173
7.2.1	Mean flow depth in explored caves (as a function of dip and catchment length)	173
7.2.2	Mean flow depth in explored caves (as a function of dip, strike and catchment length)	179
7.3	Phreatic lifts and drops	181
7.3.1	Initial phreatic drops	181
7.3.2	Terminal phreatic lifts : vauculian springs	186
7.4	Prediction of water well yields	187
7.5	Tributary junctions	188
7.6	Factors possibly modifying the initial flow net	192
7.6.1	Long vadose paths	192
7.6.2	Effects favouring shallow phreatic flow	193
7.6.3	Distortion of hydraulic conductivity ellipse by impermeable strata	194
7.6.4	The influence of topographic relief	196
7.7	Conclusion	196

Chapter 8 Flow nets in karst aquifers: response to a falling water table

8.1	Introduction	199
8.2	Longevity of flow in conduits within one tier	201
8.3	Active conduits are mostly below the water table	204
8.4	Response at initial phreatic drops and terminal phreatic lifts	206
8.5	Hydraulics of existing and developing conduits	208
8.6	There may be flooded conduits and low-gradient streams in the vadose zone	216
8.7	Cave tiers are equidistant	221
8.7.1	Previous studies	221
8.7.2	The endogenetic model	222
8.8	Hydrochemical characteristics of the flow field beneath active conduits	224
8.8.1	Expected hydrochemical characteristics	224
8.8.2	Thermal karstic springs	225
8.8.3	Sulphate / bicarbonate ratios	229
8.8.4	Implications of high sulphate values in the deep flow net	229
8.9	Conclusions	233

Chapter 9 Catchment delineation of the springs at Crowsnest Pass

9.1	Problems of catchment delineation in karst	235
9.1.1	Introduction	235
9.1.2	Groundwater boundaries do not coincide with topographic boundaries	235
9.1.3	Flows may branch to two or more springs	237
9.1.4	Flow route may vary with stage	237
9.1.5	There may be unmonitored underflow	237
9.1.6	A catchment may have non-contiguous components	238
9.1.7	Conclusion: do springs have definable catchments?	238
9.2	Catchment area model for Crowsnest Pass	239
9.2.1	Model design and validation	239
9.2.2	Model use	243
9.3	Underflow and overflow regimes	250
9.3.1	Underflow and overflow in surface creeks	250
9.3.2	Overflow springs	254
9.3.3	Ptolemy Spring: a full-flow regime	254
9.3.4	Crowsnest Spring: an underflow regime	257
	Identification as an underflow regime	257
	Quantification of the underflow component	262
9.3.5	Sublacustrine Springs: an underflow regime	264
9.4	Catchment altitude	266
9.4.1	Freeze-thaw effects	268
9.4.2	Spring temperatures	268
9.4.3	Solute concentrations	270
9.4.4	Isotopes	273
9.5	Geology of catchments	273
9.6	Definition of catchments at Crowsnest Pass	273

Chapter 10 Karst groundwater flow at Crowsnest Pass

10.1	Karst spring hydraulics	277
10.1.1	Discharge	277
10.1.2	Velocity	279
10.1.3	Area of conduits	279
10.1.4	Hydraulic gradients	281
10.1.5	Friction factors	282
10.2	Aquifer characterisation from paleohydrology	282
10.3	Aquifer characterisation from hydrochemical evidence	283
10.4	Aquifer characterisation from the karst flow model	288
10.5	Reasons for the contrasting regimes of Ptolemy Spring and Crowsnest Spring	289
10.6	Flow vectors in the Flathead and High Rock Ranges	289

Chapter 11 Karst groundwater flow in the Rocky Mountains

11.1	Introduction	293
11.2	Hydraulics	293
11.2.1	Maligne karst	293
11.2.2	Castleguard Cave	295
11.2.3	Conclusions	301
11.3	The water balance	301
11.4	Surface runoff from Rocky Mountains karst	302
11.5	Aggraded karst springs	303
11.6	Regional strike-oriented underflow in the Rocky Mountains	304
11.7	Regional dip-oriented flow in the southern Rocky Mountains	307
11.8	Regional flow to thermal springs in the northern Rocky Mountains	311
11.9	Conclusions	314

Chapter 12 Karst groundwater as a geomorphic agent

12.1	The vertical distribution of solutional erosion	317
12.2	Erosion processes and fluxes in the Rocky Mountains	322
12.3	Erosional fluxes in the Ptolemy catchment	324
12.3.1	Measurements	324
12.3.2	The altitudinal variation in erosional fluxes	329
12.4	Conclusion	330

Chapter 13 Karst hydrogeology: a new paradigm

13.1	Introduction	331
13.2	Hydrochemistry	331
13.3	Hydraulics	332
13.4	Discharge and drainage	333
13.5	Morphometric analysis of conduits and conduit flow	334
13.6	A comprehensive model of karst groundwater flow	334

Chapter 14 Conclusions

References

FIGURES

1.1	Carbonate outcrops and important karst springs of the Rocky Mountains	2
2.1	Structural and lithological relationships of the carbonate Palliser and Banff-Rundle aquifers of the Flathead and High Rock Ranges	14
2.2	Carbonate outcrops and topographic blocks of the Flathead and High Rock Ranges with groundwater flow directions, assuming a planar piezometric surface with a gradient of ≤ 0.005	16
2.3	Geological sections across the Flathead and High Rock Ranges	17
2.4	Outcrops of the Palliser and Banff-Rundle aquifers in the Flathead and southern High Rock Ranges, with areas of enclosed drainage	19
2.5	Caves of the Ptolemy and Andy Good plateaux, with dye injection points	23
2.6	Topographical subdivisions of the Flathead and High Rock Ranges, with altitude of limestone contacts and hypothetical groundwater gradients	25
3.1	Daily temperatures at Sentinel from September 1985 to August 1986	34
3.2	Measurement sites in the vicinity of Crowsnest Pass, showing dye trace routes to Ptolemy Spring and Crowsnest Spring	36-37
3.3	Measurement sites in the Flathead and High Rock Ranges	38
3.4	Stage-discharge relationships for Ptolemy Spring, Crowsnest Spring, Crowsnest River and Ptolemy Creek	39
3.5	Daily discharge of Ptolemy Spring, Crowsnest Spring and Sublacustrine Springs, September 1985 to August 1986	44
3.6	Daily discharge of Crowsnest River at the outlet of Crowsnest Lake and at Frank	45
3.7	Monthly discharge of the springs and creeks at Crowsnest Pass as a percentage of the discharge in Crowsnest River at the outlet of Crowsnest Lake, from August 1985 to September 1986	45

3.8	Instantaneous discharge of Crowsnest Spring from May 1st to September 1st 1986	46
3.9	Instantaneous discharge of Ptolemy Spring from May 1st to September 1st 1986	46
3.10	Instantaneous discharge of Ptolemy Creek from May 1st to September 1st 1986	47
3.11	Dye velocities to Ptolemy Spring and Crowsnest Spring	51
3.12	Total hardness at Ptolemy Spring and Crowsnest Spring, September 1985 to August 1986	54
3.13	Mean residence time of Crowsnest Lake, 1985-1986	54
3.14	Alkaline and non-alkaline hardness concentrations at Ptolemy Spring and Crowsnest Spring, April 15 to September 22, 1986	59
3.15	Aggressiveness of spring and creek waters at Crowsnest Pass	59
3.16	Concentrations of 16 elements in precipitation and in springs at Crowsnest Pass	65
3.17	Concentrations of nine elements in Ptolemy Spring, August 21-26, 1986	65
3.18	$\delta^{18}\text{O}$ and δD measurements from Ptolemy Spring, August 10-11, 1985	68
4.1	Recession exponents as an indicator of aquifer boundary conditions ..	79
4.2	Ratio of maximum discharge to mean discharge at Ljubljana springs	84
4.3	Cross-section of the Sierra Madre Oriental and El Abra Ranges, Mexico, at latitude 22°N , showing deep sulphate and shallow bicarbonate flow vectors (after Fish, 1977)	86
4.4	Relationship between coefficient of variation of hardness and proportion of concentrated recharge to karst aquifers	91
4.5	Summer recessions of Fontestorbes, Le Baget, and Aliou springs, Ariège, France	95
	a) gauged flow assuming fixed catchments giving an annual water balance (Fontestorbes 86km^2 , Le Baget 13km^2 , Aliou 12km^2 : after Mangin, 1975)	
	b) assuming fixed catchments of 40km^2 for Fontestorbes, 13km^2 for Le Baget, and 5km^2 for Aliou, but with an underflow loss of 15km^2 for Aliou, and an underflow gain of 700ls^{-1} for Fontestorbes.	

5.1	Flow velocity in karstic terrains	101
5.2	Velocity-discharge relationships for six cave systems (Wookey and Cheddar data from Stanton and Smart, 1981; Maligne data from Smart, 1988a; Jordtulla data from Lauritzen, 1986)	112
5.3	Ratio of maximum to minimum annual discharge for 136 karst springs (data in Table 5.2)	116
5.4	Evolution of velocity and cross-section in a karst conduit	116
5.5	Crossing of flow paths in the vadose zone of Yorkshire System, Andy Good Plateau	119
5.6	Hydraulic gradients of 18 flow routes in the ten largest catchments of the Mammoth Cave area, Kentucky (USA), (data from Quinlan and Ray, 1981)	123
6.1	Conduit porosity and density in karst aquifers	133
	a) Porosity due to explored conduits in 15 caves (data given in Table 6.1)	
	b) Conduit density for the caves in Table 6.1, and for the longest 20 and deepest 20 known caves in the world (after Courbon & Chabert, 1986)	
6.2	Flow paths in an orthogonally-jointed aquifer, using joints or bed-joint intersections	137
	a) terminology (note that this diagram is extremely simplified, and that L_x/L_y is usually >1000)	
	b) relative length of flow paths as a function of χ	
6.3	Efficiency of flow paths in karst - plan patterns of 57 allogenic and 39 autogenic flow paths (data in Table 6.1)	141
	a) sinuosity	
	b) width/length ratios	
6.4	Plan patterns of the six longest explored flow paths in karst conduits ..	142
6.5	Profiles of explored phreatic loops	146-147
	a) Doux de Coly spring, France (after Isler and Magnin, 1985)	
	b) Hölloch, Switzerland (after Bögli, 1980)	
	c) Foussoubie, France (after Le Rouk, 1989)	
	d) Wookey - Swildons, England (after Ford, 1963; Drew, 1975b; Farr, 1983)	
	e) Annette - Trou de Glaz, France (after Chevalier, 1951)	

6.6	Profiles of active and fossil caves at Crowsnest Pass, showing vertical looping	148
	a) Crowsnest Spring (after Barton, 1981)	
	b) Yorkshire System, showing two important passages developed when the water table was at or above 2230m	
6.7	Relative depth below the water table of loop crests and loop bases for 19 active and fossil phreatic paths (data listed in Table 6.3)	149
6.8	Cross-section through the low potential field surrounding a developing conduit	152
6.9	Maximum possible use of bedding planes as flow routes as a function of dip and strike	157
6.10	Relative discharge of fracture flow paths in an orthogonal flow net, using the Hagen - Poiseuille equation	157
6.11	Contrasting initial flow paths in allogenic and autogenic karstic aquifers, modelled from the Hagen - Poiseuille equation	160
6.12	Possible hydraulic gradient configurations in karst	162
6.13	Two-dimensional model of Hagen-Poiseuille flow in karst aquifers ..	165
6.14	Relative times to breakthrough for proximal and distal inputs in a karst aquifer 50km in length, based on empirical length and hydraulic gradient functions	171
7.1	Strike-oriented flowlines on a single bedding plane ABCD:	175
	a) conduit development with the bedding plane providing structural control	
	b) conduit development at sparse primary tubes or bed-joint intersections	
	c) conduit development at closely spaced primary tubes or bed-joint intersections	
7.2	Mean phreatic flow depth as a function of stratal dip and aquifer length for 17 karst aquifers	180
7.3	Phreatic flow depth and the domain of thermal springs in karst as a function of aquifer length and stratal dip	180
7.4	Flow routes aslant the strike following fractures	182
7.5	Catchment area for Fontaine de Vaucluse, France (after Drogue et al., 1983, Michelot and Mudry, 1985, and Puig, 1987)	186

7.6	Measured well yield in Pennsylvania as a function of stratal dip (after Siddiqui and Parizek, 1971) compared to the predicted relationship from Equation 7.3	189
7.7	Tributary flow fields towards a target conduit, showing equipotential lines	191
7.8	Terminal phreatic lifts and drops of selected tributary passages	191
7.9	Mean phreatic flow depth for allogenic and autogenic karsts	195
8.1	Sequential development of conduits in a karst aquifer, shown soon after abandonment of conduit A in favour of conduit B	200
8.2	Tiers of cave passages in Nelfastla de Nieva, Mexico	203
8.3	Examples of underground deltas	208-209
8.4	Ratio of discharge in developing conduit C to discharge in existing conduit B for four growth rates, corresponding to capture at stages 8, 9, 10 and 11 in Figure 8.1	211
8.5	Altitude range of phreatic loops in Ogof Ffynnon Ddu I, Wales (after Smart and Christopher, 1989)	211
8.6	Contemporaneous distributary junctions (arrowed) in part of Yorkshire System when the water table was at or above 2230m	213
8.7	Phases in the life of karst conduits	214
8.8	Patterns of vertical shaft development in the vadose zone	218
8.9	The domain of karst in terms of horizontal fracture spacing between joint conduits	220
9.1	Problems of catchment delineation in karst	236
9.2	Catchments and precipitation gauge locations used for development of runoff model	240
9.3	Subcatchments used with runoff model	244-245
9.4	Model discharge for topographic catchments	246
9.5	Mean catchments in the karst of the Flathead and southern High Rock Range, derived from the runoff model	249

9.6	Examples of overflow (Kilmarnock Creek), full flow (Upper Fording River) and underflow (Crowsnest River) regimes in the Front Ranges	252
9.7	Identification of the underflow component of Kilmarnock Creek	253
	a) best fit to Upper Fording River discharge	
	b) computed groundwater and surface components of discharge	
9.8	Discharge of Ptolemy Spring and Parrish Spring in August 1985, showing the greater variation in Parrish Spring	255
9.9	Instantaneous discharge of Ptolemy Creek and Ptolemy Spring during the snowmelt peak in 1986	255
9.10	Discharge of Ptolemy Spring and Crowsnest Spring from September 1985 to September 1986, showing contrasts in exponential recessions	256
9.11	Recession coefficients for first 24 hours after major flow peaks at Ptolemy Spring and Crowsnest Spring	259
9.12	Response of Ptolemy Spring and Crowsnest Spring to precipitation events and temperature changes, January - March 1986	261
9.13	Specific discharge from Ptolemy Spring and Crowsnest Spring during the winter and summer recessions, assuming differences to be due to aquifer boundary conditions	263
9.14	Predicted surface overflow runoff in 1985-1986 from a 41 km ² Crowsnest Spring catchment assuming identical specific discharge to Ptolemy Spring	263
9.15	Calculated catchment area for Crowsnest Spring, assuming identical specific discharge to Ptolemy Spring	265
9.16	Mean monthly discharge for Sublacustrine Springs, Crowsnest Spring and Ptolemy Spring	265
9.17	Catchment areas of Sublacustrine Springs and Crowsnest Spring as a function of season, assuming the same specific discharge as Ptolemy Spring	267
9.18	Cumulative discharge of Ptolemy Spring, Crowsnest Spring, Ptolemy Creek and Sublacustrine Springs over the snowmelt season in 1986 from snowmelt and rainfall, excluding extrapolated baseflow recession from Fall 1985	269
9.19	Relationship between temperature and discharge at Crowsnest Spring. The equation line assumes two flow components: a fixed discharge component ($T=7.5^{\circ}\text{C}$), and a variable discharge component ($T=3.8^{\circ}\text{C}$)	269

9.20	Relationship between alkalinity and altitude at Crowsnest Pass	272
10.1	Mean daily discharge for Ptolemy Spring and Crowsnest Spring, and calculated discharge for Sublacustrine Springs from May to August 1986, showing karstic response of Sublacustrine Springs	278
10.2	Bimodal dye peaks at Crowsnest Spring from Phillipps Pass injections	280
10.3	Chemograph variation at Ptolemy Spring from September 1985 to September 1986	286
	a) Total hardness	
	b) Bicarbonate and sulphate	
10.4	Chemograph variation at Crowsnest Spring from September 1985 to September 1986	287
	a) Total hardness	
	b) Bicarbonate and sulphate	
10.5	Runoff vectors from the karst of the Flathead and High Rock Ranges	292
11.1	Water levels in Medicine Lake as a function of head above the flow constriction at the lake bottom (at low discharges) and above Maligne Springs (at high discharges)	296
11.2	Plan of Castleguard Cave	298
11.3	Profile of Castleguard Cave, viewed along the strike of the strata . . .	299
11.4	Discharge from Maligne Springs from September 1st 1985 to May 1st 1986, showing the nearly constant minimum discharge between February 1st and May 1st	307
11.5	Location of major thermal springs of the Rocky Mountains	312
11.6	Variation in the proportions of bicarbonate and sulphate in overflow springs (Ptolemy Spring, Parrish Spring), underflow springs (Crowsnest Spring, Sublacustrine Spring), and thermal springs (Banff, Miette, Radium and Fairmont Hot Springs) in the Southern Rocky Mountains	315
12.1	Percentage of bedrock removed by solution as a function of depth for 9 karsts	318
12.2	Model of principal solution zones in karst	323
12.3	Hypsometric curve for the Ptolemy Creek catchment	330
12.4	Geomorphological units of the Ptolemy Creek catchment	326

12.5	Specific dissolution of limestone in the Ptolemy Creek catchment as a function of altitude	327
12.6	Erosional fluxes in the Ptolemy Creek catchment as a function of altitude	327

Tables

1.1	Discharge and hydrochemistry of the principal eastward-flowing catchments in the Rocky Mountains between Crowsnest Pass and Jasper	4
2.1	Mean composition of limestones at Crowsnest Pass (after Holter, 1976)	18
2.2	Limestone blocks of the Flathead and High Rock Ranges	24
2.3	Lithology and aquifers of the Front Ranges	28
3.1	Hydrological techniques used in karst aquifer studies at Crowsnest Pass	32
3.2	Rating curve coefficients for gauged streams at Crowsnest Pass	40
3.3	Monthly discharges of the springs and creeks at Crowsnest Pass, 1985-1986	42
3.4	Altitude and discharge of the karst springs at Crowsnest Pass	43
3.5	Dye traces at Crowsnest Pass	48
3.6	Hydrochemistry of Crowsnest River at the outlet of Crowsnest Lake . .	56
3.7	Alkaline and non-alkaline hardness of spring and creek waters at Crowsnest Pass	58
3.8	Coefficient of variation for hardness values at Ptolemy Spring and Crowsnest Spring	60
3.9	Aggressiveness of autogenic karst springs and cave streams	62
3.10	$\delta^{18}\text{O}$ results from Ptolemy Spring and Crowsnest Spring	67
4.1	Discharge budget for the White Peak, England (after Edmunds, 1971, and Christopher et al., 1977)	75
4.2	Discharge characteristics for distinguishing flow types	77
4.3	Major karst springs of the Valles - San Luis Potosí Platform, Mexico (data from Fish, 1977)	87
4.4	Variations in anion concentrations in overflow and underflow springs . .	93

4.5	Characteristics of Aliou, Le Baget and Fontestorbes Springs, Ariège, France (after Mangin, 1975, 1984)	94
4.6	Baseflow recession indices for Aliou, Le Baget and Fontestorbes Springs, Ariège in 1970 (data from Mangin, 1975)	96
5.1	Conduit groundwater velocities from multiple tracer tests	107
5.2	Annual variation in discharge (Q_x/Q_n) for karst springs	108
5.3	Recovery period of cave conduits	110
5.4	Hydraulic gradients in well-karstified limestones	113
5.5	Apparent friction factors for some karst conduits	125
6.1	Conduit density and porosity in karst aquifers	132
6.2	Conduit segment lengths	136
6.3	Sinuosity of flow routes in karst	138
6.4	Crest and base depths for active and fossil phreatic loops	144
6.5	Sulphate and bicarbonate concentrations in karst springs	168
7.1	Conduit and catchment characteristics for 16 caves	177
7.2	Notable initial phreatic drops and terminal phreatic lifts	184
7.3	Phreatic flow depth for caves in narrow karst aquifers in Norway	196
8.1	Vertical spacing between cave tiers	202
8.2	Occupancy times of cave tiers based on radiometric and paleomagnetic dating and denudation rates	203
8.3	Loop amplitude and tier spacing	204
8.4	Vadose incision in major low-gradient river caves ($Q_m > 1\text{m}^3\text{s}^{-1}$, $L > 1\text{km}$)	207
8.5	Heat fluxes of thermal springs in the Peak District, England (data from Edmunds, 1971)	228
8.6	Contrasts of temperature and sulphate/bicarbonate ratios of karst springs	230

9.1	Runoff model catchments and results	242
9.2	Areas of karst catchments draining the Flathead and High Rock Ranges	248
9.3	Discharge characteristics of the principal springs at Crowsnest Pass	256
9.4	Karst catchment areas at Crowsnest Pass	274
11.1	Major gauged karst springs of the Rocky Mountains	294
11.2	Winter discharge to some major transverse valleys in the Rocky Mountains	305
11.3	Regional hydraulic gradients from Rocky Mountain valleys east to the Prairies and west to the Rocky Mountain Trench	310
11.4	Heat fluxes of thermal and mineral springs of the Rocky Mountains ..	313
12.1	Total hardness measurements at Waitomo, New Zealand (after Gunn, 1981b)	320
12.2	Erosion processes studied in the Ptolemy catchment	325
12.3	Sediment fluxes in high mountain basins	328